

QUEENSLAND CLIMATE ASSESSMENT REPORT

Current climate and seasonal outlook information to support sustainable grazing land, water, catchment and environmental management in Queensland.

SEASONAL CONDITIONS SUMMARY

FINDINGS AS AT 13 FEBRUARY 2026

- Twelve-month rainfall to the end of January 2026, averaged over the main grazing lands of Queensland, was 43 per cent above the long-term average (1890 to 2020).
- Rainfall over the last three-, six- and 12-month periods has been well-above average to extremely high over the northern half of the State. Percentiles were in the average to well-below average range in parts of southern and southeast Queensland.
- Sea surface temperatures in the tropical Pacific indicate weakening La Niña conditions. All global models observed by BoM indicate continued warming in the tropical Pacific through to late autumn.
- As at 1 November 2025, the Science Division of DETSI considers that the probability of exceeding median summer (November 2025 to March 2026) rainfall for much of Queensland is currently in the near-average to above-average range.

IN THIS ISSUE

(click item to go to section).

- **Current Conditions - Relative to Historical Records**
 - Historical Rainfall Variability and Drought Situation
 - Rainfall and Pasture Growth Update
 - Season-to-Date Rainfall
- **Climate Drivers**
 - El Niño-Southern Oscillation (ENSO).
 - Madden-Julian Oscillation (MJO).
- **Seasonal Rainfall Outlooks**
 - Long-Lead Summer (Nov to Mar) Rainfall Outlook based on SPOTA-1
 - Three-month Rainfall Outlooks based on dynamical models
 - Three-month Rainfall Outlook based on the SOI Phase system
- **Pasture Outlook**
 - AussieGRASS three-month Pasture Growth Outlook

Historical Rainfall Variability and Drought Situation

Figure 1 shows 12-month rainfall averaged over the region of Queensland shown on the grey map (top right). This region carries over 80 per cent of the State’s cattle and sheep herd and covers only 60 per cent of Queensland’s land area. In each graph, rainfall is expressed as a percentage difference from the long-term (1890 to 2020) average. The upper graph shows the running 12-month rainfall anomaly since 1890, indicating the long-term history of wet and dry periods over this region. The lower graph focuses on changes since 2010 and shows, at selected times, those parts of Queensland drought-declared under State Government processes. Twelve-month rainfall to the end of January 2026, averaged over the main grazing lands of Queensland, was 43 per cent above the long-term average (1890 to 2020). Queensland currently has [no local government areas which are officially drought-declared](#).




Fig.1: Historical 12-month rainfall anomalies
 (Rainfall data source: <https://www.longpaddock.qld.gov.au/silo/>)

Rainfall and Pasture Growth Update

Figures 2 and 3 are products of the [AussieGRASS Environmental Calculator](https://www.longpaddock.qld.gov.au/aussiegrass) (Carter et al. 2000). Figure 2 shows rainfall over the last three-, six- and 12-month periods (up to January 31, 2026) ranked against historical values. Heavy rainfall over northern and central Queensland was linked to active wet season troughs associated with tropical cyclone Koji that made landfall near Ayr as a Category 1 system on 11 January, causing widespread flooding. Rainfall over the last three-, six- and 12-month periods has been generally well-above average to extremely high over northern half of the State. Percentiles were in the average to well-below average range in parts of southern and southeast Queensland.




Fig. 2: Rainfall percentiles (base period 1890 to current)
(Source: <https://www.longpaddock.qld.gov.au/aussiegrass/>)

Figure 3 shows calculated pasture growth percentiles over the last three-, six-, and 12-month periods. The pattern of calculated pasture growth percentiles over the last three-, six-, and 12-month periods has broad similarities for much of the state to that of rainfall percentiles (above), ranging from extremely low to extremely high depending on region and length of period considered.




Fig. 3: Pasture growth percentiles (base period 1957 to current)
White shading indicates areas that normally have "seasonally low" pasture growth.
(Source: <https://www.longpaddock.qld.gov.au/aussiegrass/>)

Season-to-Date Rainfall versus November to March historical records

Rainfall for November to January 2026 period, averaged across Queensland's major grazing lands region (grey region on the map in Fig.4), totalled 300 mm, and is above the long-term median for November to March period. The season is slightly lower than the 90th percentile and higher than the same period last year for November and January 2025.




Fig. 4: November 2025 to March 2026 season accumulated rainfall as at 1 February 2026, compared to historical values.
Percentiles based on season-to-date rainfall at the end of each month since 1 November (based on data from 1890 to 2024).
Data source: <https://www.longpaddock.qld.gov.au/silo/>

El Niño-Southern Oscillation (ENSO)

The most closely monitored driver of Queensland rainfall is the El Niño-Southern Oscillation (ENSO) phenomenon. Climate scientists monitor several ENSO indices, including the atmospheric [Southern Oscillation Index](#) (SOI) and sea-surface temperature (SST) anomalies in the central equatorial Pacific Ocean.

Rolling three-month average values of the SOI and central equatorial Pacific (Niño 3.4 region) SST anomalies since 2009 are shown in Figure 5. The most recent three-month (November to January) values of both the SOI (+6.8) and Niño 3.4 SST anomaly (-0.61°C) indicate continuing La Niña conditions (during La Niña events three-month SOI values tend to remain above +5 and Niño 3.4 SST anomalies below -0.50°C). However, La Niña and El Niño events tend to break down over autumn, and models are projecting warming in the equatorial Pacific, which DETSI would consider to be consistent with a return to ENSO-neutral conditions over coming months.




Fig. 5: El Niño Southern Oscillation (ENSO) Indices
 SST source: www.cpc.ncep.noaa.gov/data/indices
 (Niño3.4 SST, monthly OISST.v2.1 (1991 to 2020 base period))
 SOI source: www.longpaddock.qld.gov.au/soi/soi-data-files
 (monthly SOI 1887-1989 base period)





Fig. 6: Pacific Ocean sea-surface temperature anomalies (°C)
 for 11 January to 7 February 2026 based on NOAA OISST.v3.1 SST data (1991 to 2020 based period) Map source: <https://psl.noaa.gov/map/clim/sst.shtml>

Madden-Julian Oscillation (MJO)

The [Madden-Julian Oscillation \(MJO\)](#) is a broad area of tropical convection (cloud and rainfall) which tracks east from the western Indian Ocean toward the eastern Pacific Ocean. This MJO cycle tends to recur on average every 30 to 60 days (intraseasonal), distinct from El Niño and La Niña events which typically persist for three to nine months. In Queensland, the MJO has its greatest influence on rainfall during the tropical cyclone season (November to April), particularly in northern parts of the State, as its passage can enhance the strength of the northern Australian monsoon and increase the chance of tropical cyclones. For example, two tropical cyclones made landfall in Queensland during the 2023-24 season ('Kiriilly' on 25 January and 'Alfred' on 8 March), both occurred when the MJO was active in Australian-Western Pacific longitudes, unlike tropical cyclone Alfred which made landfall on 8 March 2025. The MJO is indiscernible as of 31 January 2026 and most surveyed models suggest it is likely to remain weak with little influence on Australian rainfall.



Data source: http://www.bom.gov.au/clim_data/IDCKGEM000/rmm.74toRealtime.txt

Fig. 7: Madden-Julian Oscillation intensity and phase-location

The graph shows the general location (coloured bars) and intensity (grey line) of the MJO from 1 February 2024 to 6 February 2026.
 Data source: http://www.bom.gov.au/clim_data/IDCKGEM000/rmm.74toRealtime.txt

Long-lead Summer Rainfall Outlook

(produced on 5 November, based on SPOTA-1)

The Department of the Environment, Tourism, Science and Innovation (DETSI) monitors SST anomalies in key regions of the Pacific Ocean over autumn, winter, and spring and provides objective outlooks for summer (November to March) rainfall on this basis.

The current DETSI outlook for summer rainfall in Queensland is an objective analysis of historical conditions using an index based on SST anomalies measured in key regions of the extra-tropical Pacific Ocean, which factors-in the evolving ENSO-related SST pattern in the central and southwest Pacific. Historical SST gradients across these regions, from March to October, have been related to rainfall over much of Queensland in the following summer. On this basis, as at 1 November, the Science Division of DETSI considers that the probability of exceeding median summer (November to March) rainfall is within the near- to above-normal range (50 to 80 per cent) for much of Queensland. (Figs. 8 and 9).

This is the final DETSI outlook leading up to 2025/26 summer rainfall season (November to March).

Three-month Rainfall Outlooks

based on international dynamical models

The maps in Figure 10 show rainfall probabilities for upcoming three-month periods, based on dynamical climate modelling, sourced from a range of international climate agencies. Rainfall probabilities differ for each model, on both a spatial and temporal basis, due to the different methodologies used.

For February to April period, models show normal to higher-than-normal probability of above median rainfall in different parts of coastal Queensland indicating model uncertainty.

Dynamical model outlook sources:

Australian Bureau of Meteorology <http://www.bom.gov.au/sp/sco/archive/>

UK Met Office: <http://www.metoffice.gov.uk/research/climate/seasonal-to-decadal/gpc-outlooks/glob-seas-prob>

European C3S (Copernicus Climate Change Service) multi-model: https://climate.copernicus.eu/charts/c3s_seasonal/ (C3S multi-system (7))

NOAA North American Multi-model Ensemble -USA: <https://www.cpc.ncep.noaa.gov/products/NMME/seasanom.shtml> (Global prate)

Probability of exceeding median summer rainfall for November 2025 - March 2026, as at 1 November 2025




Fig. 8: Probability of exceeding median summer rainfall (for November 2025 to March 2026) as at 1 November 2025
Produced by the DETSI SPOTA-1 system.

Summer (Nov to Mar) rainfall probabilities for Queensland regions, as at 1 November 2025




Fig. 9: Probabilities of above and below median summer rainfall (for November to March) as at 1 November 2025.

For the Queensland's major grazing region and the State as a whole, based on historical analogue year output from the DETSI SPOTA-1 system.




Fig. 10: Rainfall probabilities for February to June 2026, based on international dynamical models.

Three-month Rainfall Outlook

based on the SOI Phase system

The map in Figure 11 shows rainfall probabilities for the coming three-month period, based solely on a Rapidly Rising 'phase' of the SOI at the end of January 2026, as determined by the SOI Phase system. The current phase of the SOI ([click for more detail](#)) is based on the most recent values of the SOI (-1.5 for December and +8.0 for January 2026).

The map, which is based on 27 previous years (from 1889 to 2015) which had a 'Rapidly Rising' phase of the SOI at the end of January 2026, indicates a generally 30 to 60 per cent probability of exceeding median February to April 2026 rainfall across much of Queensland.




Fig. 11: Probability of exceeding median rainfall for February to April 2026 based on a 'Rapidly Rising' phase of the SOI at the end of January 2026.

Pasture Outlook

Figures 12 and 13 are products of the [AussieGRASS Environmental Calculator](#). Calculated pasture ground cover at the end of January 2026 (Fig. 12) ranged from average to well-above average in most parts, except for parts of southwest Queensland with below average ground cover. Figure 13 shows that the probability of exceeding median pasture growth over February to April 2026 is above-normal for most parts of Queensland, except for southwest and southern parts of the State with below-average ground cover. This calculation takes into account both current conditions (e.g., soil water content) and rainfall probabilities for the next three-month period based on [SOI Phase analogue years](#).




Fig. 12:
Calculated pasture cover at 31 January 2026.

Fig. 13:
Probability of exceeding median pasture growth for Feb to Apr 2026 based on a 'Rapidly Rising' phase of the SOI at the end of January 2026.

REFERENCES AND RESOURCES

References:

Carter, J.O. *et al.* (2000). Aussie GRASS: Australian Grassland and Rangeland Assessment by Spatial Simulation. In 'Applications of seasonal climate forecasting in agricultural and natural ecosystems - the Australian experience.' (Eds. G. Hammer, N. Nicholls and C. Mitchell). Kluwer Academic Press, Netherlands, pp. 329-349.

Day, K.A. *et al.* (2001). 'Queensland summer rainfall in relation to sea-surface temperature in the previous autumn (March) and spring (October).' Seasonal Pacific Ocean Temperature Analysis - 1 (SPOTA-1), Report 7, 2001/2. Issued 09/11/2001 on <http://www.LongPaddock.qld.gov.au>. Queensland Department of Natural Resources and Mines, Brisbane.

Day, K.A. and McKeon, G.M. (2018). An Index of Summer Rainfall for Queensland's Grazing Lands. *Journal of Applied Meteorology and Climatology*, **57**, 1623-41. <https://doi.org/10.1175/JAMC-D-17-0148.1>

McKeon, G. *et al.* (2004). 'Pasture Degradation and Recovery in Australia's Rangelands: Learning from History.' Queensland Department of Natural Resources, Mines, and Energy, Brisbane, Australia.

Stone, R. and Auliciems, A. (1992), SOI phase relationships with rainfall in eastern Australia. *Int. J. Climatol.*, **12**: 625-636. <https://doi.org/10.1002/joc.3370120608>

Stone, R., Hammer, G. & Marcussen, T. Prediction of global rainfall probabilities using phases of the Southern Oscillation Index. *Nature* **384**, 252–255 (1996). <https://doi.org/10.1038/384252a0>

Additional resources:

For further information on AussieGRASS, seasonal climate outlooks (including the SOI Phase scheme and the SPOTA-1 scheme) and Queensland drought declarations, visit the Long Paddock website, <https://www.longpaddock.qld.gov.au>.

MJO description: <https://www.climate.gov/news-features/blogs/enso/what-mjo-and-why-do-we-care>

General Disclaimer:

Information contained in this report is provided as general advice and the maps are designed to provide information relevant at a regional (shire) scale. More detailed information is required when making specific property scale management decisions. The State of Queensland, as represented by the Department of the Environment, Tourism, Science and Innovation, gives no warranty in relation to the data including without limitation, accuracy, reliability, completeness or fitness for a particular purpose.

It should be noted that seasonal outlooks are probabilistic and not deterministic in nature. For example, if an outlook indicates that there is an 80 per cent probability of exceeding median rainfall, should be interpreted as also meaning that there is a 20 per cent probability of rainfall being below the long-term median. In cases where outcomes with a high probability may be more likely, this does not mean that less probable events will not occur in any given year. Users should also consider sourcing information related to the historical track record of any outlook scheme, and such information is becoming increasingly available.

Contact: Simba Chidzambwa, Water Planning Science; Phone: 07 3170 5499; Email: simba.chidzambwa@detsi.qld.gov.au